

Preliminary Studies on the Early Life History of the Queen Conch, *Strombus gigas*, in the Exuma Cays, Bahamas

ROBERT I. WICKLUND, LUCINDA J. HEPP
& GERALDINE A. WENZ

*Caribbean Marine Research Center, Lee Stocking Island,
Exuma Cays, Bahamas,
100 East 17th Street, Riviera Beach, FL 33404*

ABSTRACT

The Caribbean Marine Research Center has initiated a study on the life history of the queen conch *Strombus gigas* in the Exuma Cays, Bahamas. Nursery grounds of *S. gigas* in the southern Exumas are described. Surveys of a major nursery area conducted from October through November 1984 resulted in a count of 570,000 juvenile 0+ and 1+-year-class conch within a total area of 386,000 m². Average density of the conch within the area was 1.48/m². Average density within grass beds and adjacent sand shoals was 1.72/m² and 0.76/m², respectively.

Modes of length-frequency curves were 8.0 cm for May 1985 and 11.0 cm for February 1986.

Seasonal distribution and behavior of deepwater adults are also described.

INTRODUCTION

The queen conch (*Strombus gigas*) is one of the most important marine species in the Caribbean region. It has been second only to finfish as a harvested resource for the past century (Brownell and Stevely, 1981). Its popularity as a food source, however, has resulted in depletion of the species in southern Florida, the Bahamas and most of the Caribbean and Bermuda. The species has been so depleted in Florida that the State has closed the fishery completely. In the Exuma Cays, Bahamas, adult queen conch are no longer found in shallow waters where they were once abundant (Richard Ellis, pers. comm.). Most adult conch in this area are located in relatively deep channels (>6 m) and offshore, out of reach of most breath-hold divers.

A paucity of data concerning the life history of the queen conch existed until recently (Berg, 1976; Brownell *et al.*, 1977; Hesse, 1979; Jory, 1982; Ballantine and Appeldoorn, 1983; Wood and Olson, 1983; Appeldoorn, 1985; Iversen *et al.*, 1986; Iversen *et al.*, 1987). The recent development of hatchery techniques has provided a better understanding of the early life stages of *S. gigas* (Brownell *et al.*, 1977; Iversen, 1983; Siddall, 1983; Appeldoorn and Sanders, 1984).

Randall (1964) made some early observations concerning predation on juvenile and adult queen conch and their migration. Recent studies suggest predation to be the most important factor in juvenile queen conch mortality (Jory, 1982; Iversen *et al.*, 1986).

Relatively little is known, however, about the factors controlling recruitment, habitat selection, density dependency and survival during the first months of settlement.

In this paper we present preliminary biological information on the life history of *S. gigas* from the southern Exuma Cays, Bahamas, during 1984, 1985 and 1986.

STUDY SITE

The southern Exuma Cays lie on the southwestern edge of Exuma Sound on the Great Bahama Bank, 110 miles southeast of Nassau, Bahamas (Figure 1). The overall study area is in the vicinity of Lee Stocking Island (23° 45' N, 76° 10' W), site of the Caribbean Marine Research Center (CMRC). Most of the study sites on the western side of the Cays have strong tidal currents, particularly between islands, resulting in shifting sand banks. Temperatures range from about 21°C in February to 30°C in August and September. Salinity ranges from 36 ppt to an occasional 40 ppt. The bottom is characterized by stable subtidal sand and seagrass flats, comprised primarily of turtlegrass (*Thalassia testudium*) and high concentrations of the calcareous algae *Rhipocephalus*, *Penicillus* and *Halimeda* species. Mounds produced by the shrimp *Callinassa* are common in the same area.

The bottom is stabilized by an organic mat consisting primarily of algae, diatoms and the rhizomes of *T. testudium*. The stabilized grassflats contain ooids, pellets, grapestones and fine-grain carbonates (Kendall and Dill, 1987).

The eastern side of the Cays is characterized by a typically open ocean profile (2 m - >30 m), coral reefs, sand flats and smooth rising relic coral reef mounds.

Norman's Pond Cay (Figure 2, Site A) is characterized by an old salt production pond (about 80 acres) with an effluent channel on the western side of the Cay. Sand and beachrock lie north and south of the channel entrance. A partial dike at the entrance to the channel maintains water level in the pond about two feet above the surrounding sea during the ebbing and low tides, resulting in an outward flow most of the day. For a short period during high tide, when the level of the sea is above the dike, the flow is into the pond.

The site of the Children's Bay Cay *S. gigas* nursery area (Figure 2, Site B), is characterized by a large bed of *T. testudium*, adjacent to an unstable ooid shoal to the west and a moderately deep channel (>7 m) to the east. The nursery ground is primarily in a grass bed at a depth of 3 - 5 m which extends onto a shallow sand shoal (1 - 2 m).

The Tugboat Cay site (Figure 2, Site C) extends from the Cay to about 1/2 mile northwest onto a sand shoal and 1 mile west. Its major bottom characteristic is sporadic patches of *T. testudium* and coarse sand pebbles with a mean depth of about 4 m. Tidal currents reach about 1 knot.




FIGURE 1. LEE STOCKING ISLAND

Figure 1. Lee Stocking Island.




FIGURE 2. STUDY SITES.

Figure 2. Study Sites.

The Leaf Cay site (Figure 2, Site D) is a small patch of *T. testudium* on the southwest side of the Cay in 4 m surrounded by deeper water. This is a minor nursery lying to the west of a major inlet.

METHODS

Standard scuba diving techniques were used to observe and measure juvenile conch. Plastic spaghetti tags were cut and tied around the spires of the conch shell.

Conch were measured along the anterior-posterior axis of the shell. Population density estimates of juvenile conch at the Children's Bay Cay site were obtained by:

1. Measuring distances and direction between buoys set on the outer perimeters of the bed with a range finder instrument.
2. Estimating area of the bed geometrically.
3. Counting and measuring conch within 20 circles each measuring 31.9 m². Fifteen of the samples were taken on grass beds and five on the adjacent sand shoal.

Surveys of a shallow-water population of *S. gigas* at Norman's Pond Cay were conducted by counting all conch in blocks, each measuring 39 m north and south alongshore and 2 m east and west seaward (Figure 3).

RESULTS AND DISCUSSION

Juvenile Queen Conch

Norman's Pond Cay

Distribution of juvenile conch (0+ and 1+ year class) at Norman's Pond Cay in July 1984 appeared to be related to the effluent from the salt pond which flows from the channel either north or south alongshore most of the day. Ambient water temperature was about 30°C and effluent temperature was about 2°C above ambient during the afternoon.

Approximately 2500 conch counted on July 18-28, 1984, were centered at the entrance to the channel and spread directly seaward about 22 m from the entrance and along the shoreline north and south (Figure 3). The conch were often exposed along the beach at low tide. This behavior has also been reported by Iversen *et al.* (1987).

By August 18th the conch started moving away from the channel entrance to the north and south. On September 9th only two conch were observed at the channel entrance, but heavy concentrations were found along the shallow shore on both sides. Ambient water temperature was 31.6°C, shoreline temperature was 31.6°C, and effluent temperature was 33.3°C.

On October 7th most conch were concentrated in two large clusters on the south side of the inlet. Waterflow from the channel at this time was northward alongshore. Ambient temperature was 26°C; effluent temperature reached 27°C

		A	B	C	D	E	F	G	H	I	J	K	L
NORTH	1	16	13	10	32	47	33						
	2	24	5	11	19	9	5						
	3	28	12	28	15	12	11						
	4	62	61	40	23	8	4	0					
	5	7	36	9	9	2	2	0					
CHANNEL ENTRANCE	6	103	81	88	47	51	45	45	53	59	37	21	0
SOUTH	7	24	26	21	27	44	15	7					
	8	42	50	36	24	29	8	2					
	9	72	68	38	7	32	7	12					
	10	86	56	47	23	17	9	2					
	11	39	64	28	37	6	3	2					
	12	47	51	34	12	1							
	13	26	38	13	6	3							
	14	50	27	11	1	1							
	15	7	1	2	0	0							

Figure 3. Distribution and number of juvenile *S. gigas* surveyed at the entrance channel to the Old Salt Pond located at Norman's Pond Cay, Bahamas, 18-24 July 1984.

by noon. This was the last observation of the Norman's Pond Cay population of conch due to an almost complete harvest by local fishermen soon after October 7th.

We presume that the association of juvenile *S. gigas* with the salt pond effluent was related to higher temperature and possibly higher nutrient content of effluent water (which was normally a murky green color), particularly during the late spring to early fall months. However, in July 1984 we observed that during the ebbing tide, when the channel water flows seaward, the conch moved away from the entrance as illustrated in Table 1.

We believe this movement of conch away from the channel during the outgoing tide was due to either the strong current coming from the channel or abnormally high temperatures of effluent water occasionally found at midday.

Table 1. Number of Conch at Norman's Pond Cay Channel Entrance and Adjacent Sites at High and Low Tide - July 1984.

	High Tide	Low Tide
North	29	95
Entrance	184	95
South	50	110

During late summer, when temperatures of the pond effluent reached 33.3°C, the conch moved away from the channel completely but remained along the shoreline.

In October 1984 westerly storms and winds brought relatively high wave action onto Norman's Pond Cay. Following this we observed that for one week many of the young *S. gigas* were partially buried or gathered around small coral heads. Presumably this behavior was in response to the high energy waves in shallow water that could wash the conch onto the beach. Only a few dead conch were found on the beach stranded after the storm. Appeldoorn (1985) reported that burial of *S. gigas* and *S. costatus* was greater during periods of turbulence in Puerto Rican waters.

Children's Bay Cay

Our pilot studies were centered around a large *S. gigas* nursery area adjacent to Children's Bay Cay just south of Lee Stocking Island during 1985 and 1986. This nursery area is characterized by *Thalassia* grass, coral rubble, stabilized sand and adjacent shifting ooid shoals. Depth ranges between 3 and 5 m and tidal currents move west and east at about 1 knot at the peak of flow.

Over a 3-year period this area remained populated with juveniles rarely exceeding 120 mm, although an occasional lipped adult was found in the nursery area. As of May 1985 no individuals measured under 60 mm. This is consistent with reports that juvenile *S. gigas* under 80 mm bury and are out of sight during the day (Randall, 1964; Appeldoorn and Ballantine, 1983).

Density of juvenile *S. gigas* in the Children's Bay Cay nursery area appeared fairly constant throughout the year. During October 29 through November 4, 1985, we counted 570,000 juvenile *S. gigas* in an area of 386,000 m².

Mean density was 1.48/m² (N=946). Density within the two distinct habitats of the nursery area was 1.72/m² (N=824) for the *Thalassia* grassbeds and 0.76/m² (N=122) on the adjacent ooid sand shoal. Comparative density measurements in the Berry Islands, Bahamas, ranged from 1.5/10m² at Little Cockroach Cay to 19.6/10m² at Bird Cay Channel (Iversen *et al.*, 1987).

With the exception of Bird Cay Channel, the densities of juvenile *S. gigas* in the Children's Bay area of the Exuma Cays are among the highest reported

anywhere. Densities of 0.001/m² and 0.9/m² have been reported for *S. gigas* populations in the Virgin Islands (Woods and Olsen, 1983) and in the Turks and Caicos (Hesse, 1979), respectively.

The size distribution of juvenile *S. gigas* in the Children's Bay Cay nursery area ranged from 6.0 - 12.5 cm in May 1985 to 7.0 - 15.0 cm in January 1986. The modes of the length-frequency curves were 8.0 cm for May 1985 and 11.0 cm for February 1986. Iversen's (1987) estimates of average length by ages suggest that the main population we measured at Children's Bay Cay was 1 year old. This agrees with age-size estimates in Puerto Rico (Berg, 1976) and Venezuela (Brownell, 1977), but is under Von Bertalanffy growth curve analyses of populations in the Virgin Islands (Berg, 1976; Brownell *et al.*, 1976) and Cuba (Alcolado, 1976).

One hundred and fifty-two juvenile conch were tagged at the Children's Bay Cay site with 48 returns including 8 captured twice. Most of the recaptures were made within 1 - 2 months of the initial tagging and all within a short distance of the site. Mean growth rate calculated from all of the returned tags was 0.12 mm/day and 0.37 cm/month. Iversen *et al.* (1987) reported growth rates from the Berry Islands as 0.44 - 1.63 cm/month for the summer and 0.18 - 0.30 cm/month for the rest of the year. Table 2 summarizes the tagging returns for this site.

We believe that the Children's Bay Cay site is important to the recruitment of conch into the southern Exuma Cays. Our observations indicated that the numbers of 1-year-old juvenile *S. gigas* remain relatively stable suggesting that new conch recruit into the area each year. The conditions that allow conch to thrive here are not understood. Obviously, food availability, good water quality, bottom sediment suitable to allow the youngest conch to bury and avoid predation, and currents that carry the pelagic veligers to nursery areas are all important to the success and survival of the juveniles. The Children's Bay Cay site offers rich *Thalassia* grassbeds that provide food for the conch and strong tidal currents that move water in and out of Exuma Sound.

Leaf Cay

A small population of juvenile *S. gigas* was found adjacent to Leaf Cay just north of Lee Stocking Island. This group was fairly isolated and warrants only a brief mention here. Measurements made from July 1985 to January 1986 showed a size distribution ranging from 8.0 - 13.5 cm, similar to that of the Children's Bay Cay population. Modes of the length-frequency curves ranged from 10.0 cm in July 1985 to 11.5 cm in January 1986.

On June 4 and 10, 1985, 97 juvenile *S. gigas* were tagged at the Leaf Cay site. Fifty-nine were recaptured in the same area, including 14 captured twice. Mean growth rates calculated from tagging data were 0.14 mm/day and 0.41

Gulf and Caribbean Fisheries Institute

Table 2. Summary of Tag Returns.

Tagging Date	Size (cm)	Children's Bay Cay Site		Recovery Date	Size (cm)	Growth mm/day
		Recovery Date	Size (cm)			
5-02-85	9.5	6-12-85	10.0			0.12
	8.5	7-10-85	9.0			0.007
	9.5	7-10-85	10.2	8-22-85	10.7	0.01/0.11
	9.5	6-12-85	9.5			0.00
	9.5	6-11-85	10.5			0.25
	8.5	6-12-85	9.0			0.12
	9.0	6-18-85	9.0			0.00
	9.5	6-11-85	9.5			0.00
	5.5	7-10-85	6.5			0.14
	9.5	6-12-85	10.5			0.24
	8.0	6-12-85	Dead			---
	10.0	8-22-85	Dead			---
	5-13-85	9.5	6-12-85	9.5	7-10-85	10.0
9.0		6-11-85	9.5			0.17
10.0		10-21-85	Dead			---
9.5		6-12-85	10.0			0.16
9.5		7-10-85	10.0			0.08
8.5		6-11-85	8.5	7-10-85	9.0	0/0.17
9.5		7-10-85	10.0			0.08
8.0		6-12-85	8.5			0.16
9.5		8-22-85	10.5			0.09
9.0		6-11-85	9.0			0.00
8.0		6-12-85	8.5			0.16
5-15-85	8.5	6-12-85	9.0			0.16
	8.5	6-11-85	9.0			0.18
	9.0	10-21-85	11.0			0.13
	6.5	10-14-85	Dead			---
	8.5	6-12-85	9.0			0.17
	8.5	6-11-85	9.0			0.17
	9.0	6-12-85	9.5	8-12-85	10.5	0.17/0.16
	9.5	6-11-85	9.5	7-10-85	10.0	0/0.17
	8.0	6-11-85	9.0			0.37
	9.5	6-11-85	9.5			0.00
	9.0	6-11-85	10.0	10-21-85	10.5	0.37/0.04
	8.5	7-10-85	9.5	8-08-85	10.0	0.18/0.17
	8.0	6-11-85	8.5			0.18
	8.5	6-12-85	9.0			0.18
8.5	6-11-85	9.0	7-10-85	9.5	0.18/0.17	
10.5	6-11-85	10.5			0.00	
8.5	9-26-85	Dead			---	

Mean growth rate = 0.12 mm/day

cm/month. Tagging returns and growths from Leaf Cay are summarized in Table 3.

Tugboat Cay

One of the largest juvenile conch populations in the vicinity of Lee Stocking Island was found west of Tugboat Cay. Although we do not have density figures for this area, it was generally observed that densities were not as high as in the Children's Bay Cay population, the Tugboat Cay nursery area being larger and the range of sizes and ages much wider. In August 1985 sizes at Tugboat Cay (N = 78) ranged from 8.4 - 14.7 cm (Figure 5). During the same month, sizes at Children's Bay Cay (N=100) ranged from 8.0 - 12.5 cm (Figure 4). A bimodal size distribution curve of juvenile *S. gigas* was evident at various times from July 1985 - January 1986 (Figure 5) due to occasional sampling of a population of smaller conch toward the eastern end of the Tugboat area. At the Tugboat Cay site conch ranged from 1 year to adult, the dominating ages being 1 and 2 years old with significant numbers of 3+ also found. Nowhere else in the area between Leaf Cay and south of Children's Bay Cay have we found significant numbers of 2 - 3-year-old conch as were found at Tugboat Cay. Considering its close proximity, the Children's Bay Cay 1 - 2-year-old conch could migrate to the Tugboat Cay site.

Adult Queen Conch

Horseshoe Reef, Lee Stocking Island.

Adult queen conch in the Exuma Cays are now only found in relatively deep water (>6 m), generally in channels with high concentrations of *Thalassia* and offshore on sand and relic coral reef outcroppings.

The most significant adult population in the Lee Stocking Island area is an offshore spawning stock found at a depth of 15 - 23 m on the Exuma Sound side of the Island and ranging in size from 17.7 - 26.8 cm. They are found mainly during late spring through early fall on carbonate sand bottom partially covered by a thin algal mat adjacent to a deepwater reef (Horseshoe Reef) running parallel to, and about 1/2 mile from, the beach. This reef, with a steep vertical profile of about 5 m, is a barrier which prevents the conch from moving further inshore. During the winter months most of the offshore group have been observed on top of relic patch reefs. These large mounds have little living hard coral but are covered by algae and soft corals. The algae most likely provide food to the conch during the winter. During August 1987 some individuals remained on the offshore mounds (Allan Stoner, University of Puerto Rico, pers. comm.). We believe that a sizable portion of the Children's Bay Cay, Tugboat Cay and other nursery conch populations in the area are recruited from the offshore spawning groups. The depth of the offshore conch population protects

Table 3. Summary of Tag Returns.

Tagging Date	Size (cm)	Leaf Cay Site		Recovery Date	Size (cm)	Growth mm/day	
		Recovery Date	Size (cm)				
6-04-85	10.5	7-08-85	10.5			0.00	
	9.5	7-08-85	10.0			0.15	
	9.5	7-08-85	10.0			0.15	
	11.5	8-09-85	12.2			0.09	
	9.7	7-08-85	10.0			0.09	
	9.7	7-08-85	9.8			0.03	
	8.5	7-08-85	9.2	12-13-85	11.0	0.20/0.11	
	10.0	7-08-85	10.3	8-09-85	10.7	0.09/0.12	
	8.5	7-08-85	9.2			0.20	
	10.0	7-08-85	10.7			0.20	
	9.5	7-08-85	10.5			0.29	
	10.0	7-08-85	10.7			0.20	
	6-10-85	11.2	7-09-85	11.2	8-15-85	11.3	0/0.02
		10.5	7-08-85	11.0			0.17
10.0		7-08-85	10.5			0.17	
8.5		7-08-85	9.0	8-09-85	9.2	0.17/0.06	
9.5		7-08-85	10.0	8-09-85	10.2	0.17/0.06	
10.0		7-08-85	10.5			0.17	
8.5		7-08-85	9.5			0.35	
9.5		7-09-85	10.0			0.17	
10.5		7-09-85	10.5			0.00	
9.0		7-08-85	9.5			0.18	
9.5		7-08-85	10.1	8-15-85	10.1	0.21/0	
10.0		7-08-85	10.5			0.17	
10.0		7-08-85	10.5			0.17	
10.5		7-08-85	11.0	8-09-85	11.3	0.17/0.09	
10.5		7-08-85	10.5			0.00	
10.0		7-08-85	10.5			0.17	
10.0		7-08-85	10.5	8-09-85	10.8	0.17/0.09	
10.0	7-09-85	10.5			0.17		
9.5	11-06-85	11.8			0.15		
11.0	7-08-85	11.3	8-15-85	12.3	0.10/0.26		
10.5	8-15-85	10.9			0.06		
9.5	7-08-85	10.0	12-13-85	Dead	0.17		
9.5	7-08-85	9.7			0.11		
11.0	7-08-85	11.6			0.21		
9.5	7-08-85	10.0			0.17		

Table 3 (continued).

Tagging Date	Size (cm)	Leaf Cay Site		Recovery Date	Size (cm)	Growth mm/day
		Recovery Date	Size (cm)			
	9.5	7-01-85	10.0	7-08-85	10.0	0.24/0
	10.2	7-08-85	10.7			0.17
	9.2	7-09-2935	9.8			0.20
	10.2	7-09-85	10.7			0.17
	9.5	7-08-85	Dead			—
	10.5	7-01-85	11.0	7-08-85	11.0	0.24/0
	9.3	7-09-85	9.5			0.17

Mean growth rate = 0.14 mm/day

them from most fishing pressure allowing their numbers to remain constant and ensuring spawning success.

Studies of conch by other investigators indicate that their spawning season ranges from March to September (Randall, 1964; D'Asaro, 1965; Brownell, 1977) and that they migrate offshore in the winter (Randall, 1964; Hesse, 1979). We observed spawning behavior in this deepwater population on the sand bottom throughout spring and summer in 1985, 1986 and 1987. One small shallow water (<5 m) spawning group was also observed near the eastern shore of Norman's Pond Cay in July 1987.

In summary, the area around Lee Stocking Island is a major nursery ground for juvenile *S. gigas* and an excellent site for research on the species. We suggest that this site be the focus of future research concentrating on the following:

1. Recruitment of young into the nursery areas,
2. Description of factors necessary for the survival of the youngest conch,
3. Predation and
4. Food availability.




Figure 4. Size frequencies of juvenile *Strombus gigas*, Children's Bay Cay site, Exuma Cays, Bahamas -- 1985-86.

Figure 4. Size frequencies of juvenile *Strombus gigas*, Children's Bay Cay site, Exuma Cays, Bahamas -- 1985-86.




Figure 5. Size frequencies of tagged juvenile *Strombus gigas*, Tugboat Cay, Exuma Cays, Bahamas, 1985-6.

ACKNOWLEDGEMENTS

This research was supported by the Office of Undersea Research, National Oceanic and Atmospheric Administration, U.S. Department of Commerce (Grant Number NA85AA-D-UR034). Further, special thanks go to Myffie Lewis, a student from New Zealand, who set up and conducted the Norman's Pond Cay survey, Gay Van Zandt for her wordprocessing and review of this paper, and Lisa Ellingson for her illustrations. We also thank Drs. Wade Watanabe and Allan Stoner for their review and comments. Finally, this work could not have been conducted without the support of the Department of Fisheries, Bahamas Ministry of Agriculture and Fisheries.

REFERENCES

- Alcolado, P.M. 1976. Crecimiento, variaciones morfológicas de la concha y algunos datos biológicos del cabo, *Strombus gigas*, L. (Molusca; Mesogastropoda). Acad. Cienc. Cuba, Ser. Oceanol. 34, 36 pp.
- Appeldoorn, R.S. and Ballantine, D.L., 1983. Field release of cultured queen conchs in Puerto Rico: implications for stock restoration. *Proc. Gulf Carib. Fish. Inst.* 35:89-98.
- Appeldoorn, R.S. and I.M. Sanders. 1984. Quantification of the density-growth relationship in hatchery-reared juvenile conchs (*Strombus gigas* Linne and *S. costatus* Linne). *J. Shellfish Res.* 4(1):63-66.
- . 1985. Growth, mortality and dispersion of juvenile laboratory-reared conchs, *Strombus gigas* and *S. costatus*, released at an offshore site. *Bull. Mar. Sci.* 37:785-793.
- Ballantine, D.L., and R.S. Appeldoorn. 1983. Queen conch culture and future prospects in Puerto Rico. *Proc. Gulf Carib. Fish. Inst.* 35:57-63.
- Berg, C.J., Jr. 1976. Growth of the queen conch, *Strombus gigas*, with a discussion of the practicality of its mariculture. *Mar. Biol.* (Berl.) 34:191-199.
- Brownell, W.N. 1977. Reproduction, laboratory culture, and growth of *Strombus gigas*, *S. costatus* and *S. pugilis* in Los Roques, Venezuela. *Bull. Mar. Sci.* 27:668-680.
- , C. J. Berg, Jr., and K. C. Haines. 1977. Fisheries and aquaculture of the conch, *Strombus gigas*, in the Caribbean. In H. B. Stewart, Jr., (editor). Cooperative investigations of the Caribbean and Adjacent Regions, Caracas, Venezuela, 12-16 July 1976, pp. 59-69. FAO Fish. Rep. 200.
- and J. M. Stevely. 1981. The biology, fisheries and management of the queen conch, *Strombus gigas*. *Mar. Fish. Rev.* 43(7):1-12.
- D'Asaro, C.N. 1965. Organogenesis, development and metamorphosis in the queen conch, *Strombus gigas*, with notes on breeding habits. *Bull. Mar. Sci.* 15:359-416.
- Hesse, K.O. 1979. Movement and migration of the queen conch, *Strombus*

-
- gigas*, in the Turks and Caicos Islands. *Bull. Mar. Sci.* 29:303-311.
- Iversen, E.S. 1983. Feasibility of increasing Bahamian conch production by mariculture. *Proc. Gulf Carib. Fish. Inst.* 35:83-88.
- , D.E. Jory, and S.P. Bannerot. 1986. Predation on queen conchs, *Strombus gigas*, in the Bahamas. *Bull. Mar. Sci.* 39:61-75.
- , E.S. Rutherford, S.P. Bannerot, and D.E. Jory. 1987. Biological data on Berry Islands (Bahamas) queen conchs, *Strombus gigas*, with mariculture and fisheries management implications. *Fish. Bull.* 85 (2):229-310.
- Jory, D.E. 1982. Predation by tulip snails, *Fasciolaria Tulipa*, on queen conchs, *Strombus gigas*. M.S. Thesis. Univ. Miami, Coral Gables, FL, 73 pp.
- Kendall, C.G. St. C., and R.F. Dill. 1987. Guidebook to the giant subtidal stromatolites and carbonate facies of Lee Stocking Island, Bahamas. Dept. Geol. University of South Carolina.
- Randall, J.E. 1964. Contributions to the biology of the queen conch, *Strombus gigas*. *Bull. Mar. Fish. Rev.* 47(4):1-10.
- Siddall, S.E. 1983. Biological and economic outlook for hatchery production of juvenile queen conch. *Proc. Gulf Carib. Fish. Inst.* 35:46-52.
- Wood, R.S., and D.A. Olsen. 1983. Application of biological knowledge to the management of the Virgin Islands conch fishery. *Proc. Gulf Carib. Fish. Inst.* 35:112-121.