

# **Recruitment of Spiny Lobsters, *Panulirus argus*, to Submerged Sea Cages off Puerto Rico, and its Implication for the Development of an Aquaculture Operation**

MEGAN DAVIS<sup>1</sup>, BRIAN O'HANLON<sup>2</sup>, JOSÉ RIVERA<sup>3</sup>,  
JERRY CORSAUT<sup>1</sup>, TONY WADLEY<sup>1</sup>, LEROY CRESWELL<sup>4</sup>,  
JOSEPH AYVAZIAN<sup>2</sup>, and DANIEL BENETTI<sup>5</sup>

<sup>1</sup>*Harbor Branch Oceanographic Institution  
5600 US 1 North,  
Fort Pierce, Florida 34946 USA*

<sup>2</sup>*Snapperfarm, Inc.  
P.O. Box 685*

*Culebra, Puerto Rico 00775*

<sup>3</sup>*U.S. National Marine Fisheries Service  
HC-01 Box 1736*

*Boqueron, Puerto Rico 00622-9704*

<sup>4</sup>*Florida Sea Grant College Program  
8400 Picos Road, Suite 101*

*Fort Pierce, Florida 34945-3045 USA*

<sup>5</sup>*Rosenstiel School of Marine and Atmospheric Science  
University of Miami  
4600 Rickenbacker Causeway  
Miami, Florida 33149 USA*

## **ABSTRACT**

Spiny lobster, *Panulirus argus*, is an important fisheries species in Florida and the Caribbean region. The high value of this species and the limited extent of their wild fisheries make them an ideal candidate for commercial culture. One of the major constraints to spiny lobster culture is that the aquaculturist must rely on the steady supply of post larvae (pueruli) from the wild for nursery and growout. The focus of this research project was to advance the culture of spiny lobster through novel pueruli collection methods. Recruitment of 1000's of spiny lobster pueruli to submerged fish sea cages in Puerto Rico was first observed by Snapperfarm, Inc. in spring of 2003. With funds from NOAA SBIR Phase I, studies were conducted from July 2003 to January 2004 to determine the feasibility of collecting spiny lobsters from sea cages for growout. Results showed that year-round collection of pueruli from submerged sea cages is feasible, with the highest collection occurring in the spring and near the new moon phase. Newly settled pueruli and pigmented post larvae were observed during each month of the study. Over 400 juvenile lobsters were collected from the submerged sea cages; 40 were placed in a growout study and the rest were relocated to a nearby marine reserve. The findings from this study indicate that collection of lobster pueruli and juveniles

---

---

from sea cages for growout is technically feasible and has potential to be developed into a commercial venture.

KEY WORDS: Pueruli, recruitment, spiny lobster

### **Reclutamiento de la Langosta Espinosa, *Panulirus argus*, a Jaulas Sumergidas en Puerto Rico, y su Implicacion para el Desarrollo de una Industria de Acuicultura**

La langosta espinosa, *Panulirus argus*, es una especie importante en las pesquerias de la Florida y la region del Caribe. El alto valor commercial de esta especie y la distribucion geografica limitada de su pesqueria la hacen un candidato ideal para cultivo a nivel comercial. Uno de los mayores factores limitantes al cultivo de la langosta espinosa es que el acuicultor depende en el continuo abastecimiento de post larvas (pueruli) silvestres para establecer viveros y crianza a la etapa de adulto.

El enfoque de este proyecto de investigacion es la innovacion de metodos de el cultivo de los puerulus de la langosta espinosa. Reclutamiento de miles de puerulus de la langosta espinosa a jaulas sumergidas en Puerto Rico fue observado por Snapperfarm, Inc. en la primavera del 2003. Con fondos de NOAA SBIR (Primera fase), estudios fueron realizados entre Julio 2003-Enero 2004 para determinar la posibilidad de coleccionar la langosta espinosa de las jaulas sumergidas para la crianza.

Los resultados demuestran que los pueruli se pueden coleccionar a traves del ano de las jaulas sumergidas, con las mayores cantidades obteniendose en la primavera cerca de la fase de luna nueva. Los pueruli recién llegados y post larvas pigmentadas fueron observadas cada mes del estudio. Mas de 400 langostas juveniles fueron coleccionadas de las jaulas sumergidas; 40 fueron asignadas a un estudio de crecimiento y el resto fueron transferidas a una reserva marina cercana al lugar del estudio. Los resultados de este estudio demuestran que la coleccion de pueruli y juveniles desde las jaulas sumergibles es tecnicamente posible y que hay potencial para desarrollar una industria a nivel comercial.

PALABRAS CLAVES: Acuicultura, langosta espinosa, coleccion de pueruli

#### INTRODUCTION

The world market for spiny lobsters is approximately 80,000 metric tons annually and is supplied almost entirely by wild fisheries (FAO, 2005). The spiny lobster, *Panulirus argus*, is found throughout the Gulf and Caribbean and forms the largest commercial fishery for spiny lobsters in the world at around 38,000 metric tons. The high value of spiny lobsters and the limited extent of their wild fisheries are generating an increasing worldwide interest in aquaculture (Jeffs and Hooker 2000). The commercial aquaculture of spiny lobsters is estimated at 3,000 tons per year and is being cultured in a number of countries including Japan, Australia, Taiwan, and New Zealand with the vast majority

---

---

grown in Vietnam (Jeffs and Hooker 2000). This spiny lobster aquaculture is based on the collection of post-larva and early juveniles from the wild.

The prospects for farming the Caribbean spiny lobster, *P. argus*, profitably are higher than for temperate spiny lobster species due to the greater availability of wild seed lobsters and a faster growth rate (Booth and Kittaka 2000, Jeffs and Davis 2003). At this time, the spiny lobster, *P. argus*, is not commercially cultured, however, a great deal of research has been accomplished on this species and similar species that is directly relevant to the development of aquaculture of *P. argus* (Lellis 1991, Jeffs and Hooker 2000).

One of the major constraints to spiny lobster culture is obtaining a steady supply of post larvae for nursery and growout. Spiny lobsters have been successfully mated and spawned in captivity. However, researchers have had difficulty rearing large numbers of phyllosoma larvae through the many stages of the year long complex cycle. An alternative is to bypass the larviculture stage and collect the pueruli from the wild as they drift to nearshore habitats. There has been good success with the collection of wild spiny lobster pueruli on floating artificial habitats (Witham collectors), in crevice collectors, or large plankton nets (Witham et al. 1964, Calinski and Lyons 1983, Reid et al. 1991, Gutierrez-Carbonell et al. 1992, Field and Butler 1994, Phillips and Booth, 1994). The transparent pueruli settle on the collectors during the new moon to the quarter moon phase. Currents, tides, wind, and season all play a role in settlement and collection numbers. In addition to lunar cycles, wind, tides, hydrological features and seasonality influence the timing and magnitude of lobster recruitment to artificial habitats.

Collection of pueruli from the natural environments will not put pressure on the fisheries stocks. There are indications from pueruli collection studies that collectors only remove a small fraction of the number of pueruli in the water column and that a very large number of pueruli are lost mainly due to predation or lack of appropriate settlement habitat (Ryther et al. 1988, Butler and Herrnkind 1989). In nature it is estimated that only 4% of post larvae survive to juvenile stage (Forcucci et al. 1994). Under controlled culture conditions survival could be as high as 50 - 75% during this stage (Lellis 1991).

Snapperfarm, Inc. has two 3,000 cubic meter sea cages used to culture marine fish off the coast of Culebra, Puerto Rico. This structure is located in 28 meters of water depth. Over the course of the first six months after deployment, the cage became covered in a fine layer of macroalgae and diatoms. In the spring of 2003, 1000's of spiny lobster pueruli recruited to these submerged sea cages. These juvenile lobsters were the first indication that these large sea cages are ideal settlement collectors for spiny lobster pueruli. With funds from NOAA SBIR Phase I, Snapperfarm, Inc. and researchers from Harbor Branch Oceanographic Institution, University of Miami - RSMAS, Florida Sea Grant Extension, and NMFS conducted studies from July 2003 - January 2004 to advance the culture of spiny lobster through novel pueruli collection methods.

---

---

**OBJECTIVES:**

The main goals of the project were to determine methods to collect spiny lobster pueruli that recruited to offshore submerged sea cages in Puerto Rico, and to assess the commercial viability of a spiny lobster mariculture operation in association with fish sea cages.

**METHODS AND RESULTS**

The study was conducted by Snapperfarm, Inc. and collaborators in Culebra, Puerto Rico. Snapperfarm has two SeaStation™ 3,000 cubic meter offshore cages off the coast of Culebra, Puerto Rico. Each cage is 15 m deep, 25 m in diameter and has a total surface area of 1,115 square meters. The cages are submerged in a water depth of 28 meters with the top 8 m below the sea surface. The site is fully exposed to the Atlantic Ocean to the northwest and the Caribbean Sea to the southeast. The site receives strong, open ocean currents (0.5 knots).

Scuba diving surveys of the sea cages for spiny lobster pueruli were conducted during each month of the study (July 2003 – January 2004) and showed that the majority of the lobsters recruited to the sea cages on the new moon and during the week following the new moon. Low numbers of pueruli were visually found during transect surveys of the sea cages, due to the small size of the pueruli (6 - 7 mm CL) and their cryptic nature (e.g., clear to slightly pigmented). The most number of pueruli (total of 14) were seen while night diving with lights on a small section of the sea cage in August. Pueruli were also observed in the daytime during each month of the study (up to five pueruli per month). The pueruli were found near the chimney top of the sea cage, in the folds of the harvest net, and under clumps of red algae, *Laurencia* sp., which is known to trigger lobster settlement. This is the first time that lobsters of this size have been observed in the wild on large submerged sea cages.

Collection devices, such as PVC pipes and window screen pinwheels, were secured to the sea cages in an experimental array. Lobsters did not recruit to the PVC pipes, but were attracted to the screen pinwheels. Up to five pueruli and juveniles were found in the screen pinwheel collectors each month (October 2003 – January 2004). The best results were when the pinwheels were placed on the sea cages one week prior to the new moon for fouling. However, these collection devices (PVC pipes and pinwheels) were not ideal for pueruli and juvenile lobster collection, because of the crabs that also recruited to the collection devices.

Due to the cryptic nature of the pueruli, an alternative collection method used during the study included removing spiny lobster juveniles off the sea cages after they had gone through several molts (two months after settlement). Over 400 juvenile lobsters (two to eight months old) were collected from the two sea cages during the study. The majority were placed in a nearby marine reserve, however, 40 were removed for a preliminary grow out study. They ranged in size from 3 – 6 cm CL (five to eight months in age). The lobsters were placed inside a 55-gallon plastic barrel that had holes drilled in the sides and shelter inside. The barrel was secured to the sea cage mooring on bottom of the ocean (28 m deep). The lobsters were fed every five to seven days with

a 7 – 9 kg cobia fish that had recently died in the sea cage. Lobster survival over the three month study period was 60%, however, growth was slow over the three months due to lack of adequate feed variety.

#### DISCUSSION

There is every indication from this study that lobster pueruli and juveniles can be collected year round from submerged sea cages in Puerto Rico. Due to the cryptic behavior and clear color of the pueruli, it is not surprising that it was difficult to obtain an accurate count of the number of pueruli during the surveys. The results from this study show that year round collection of lobster pueruli and juveniles from sea cages is technically feasible and has potential to be developed into a pilot-scale venture. This is consistent with the purpose of NOAA SBIR Phase I to determine the technical feasibility of the proposed research. Recruitment of lobsters to the fish sea cages will provide Snapper-farm, Inc. with an added revenue source to support the fish farming operation. The company was awarded a NOAA SBIR Phase II to continue the research and development of pueruli collection on submerged sea cages and submerged cage growout of juveniles for market.

This is Harbor Branch Oceanographic Institution Contribution Number 1567.

#### LITERATURE CITED

- Butler, M.J. and W.F. Herrnkind. 1989. Are artificial "Witham" surface collectors adequate indicators of Caribbean spiny lobster, *Panulirus argus*, recruitment. *Proceedings of the Gulf and Caribbean Fisheries Institute* **42**:135-136.
- Booth, J.D. and J. Kittaka. 2000. Spiny lobster growout. Pages 556-585 in: B.F. Phillips and J. Kittaka (eds.). *Spiny Lobster Management*. Fishing News Books, Oxford, United Kingdom.
- Calinski, M.D. and W.G. Lyons. 1983. Swimming behavior of the puerulus of spiny lobster *Panulirus argus* (Latreille, 180) (Crustacea: Plainuridae). *J. Crustacean Biology* **3**:329-335.
- FAO (Food and Agriculture Organization of the United Nations). 2005. Fishery statistics, catches and landings. FAO yearbook-Fisheries. Food and Agriculture Organization of the United Nations, Rome, Italy.
- Field, J.M. and M. Butler. 1994. The influence of temperature, salinity, and postlarval transport on the distribution of juvenile spiny lobsters, (Latreille, 1804), Florida Bay. *Crustaceana* **67**:26-45.
- Forcucci, D., M.J. Butler, and J.H. Hunt. 1994. Population dynamics of juvenile Caribbean spiny lobster, *Panulirus argus* in Florida Bay, Florida. *Bulletin of Marine Science* **54**:805-818.
- Gutierrez-Carbonell, D., J. Simonin-Diaz, and P. Briones-Fourzan. 1992. A simple collector for postlarvae of the spiny lobster *Panulirus argus*. *Proceedings of the Gulf and Caribbean Fisheries Institute* **41**:516-527.

- 
- 
- Jeffs, A.G. and S. Hooker. 2000. Economic feasibility of aquaculture of spiny lobsters *Jasus edwardsii* in temperate waters. *Journal of the World Aquaculture Society* **31**:30-41.
- Jeffs, A.G. and M. Davis. 2003. An assessment of the aquaculture potential of the Caribbean Spiny Lobster, *Panulirus argus*. *Proceedings of the Gulf and Caribbean Fisheries Institute* **54**:413-426.
- Lellis, W. 1991. Spiny lobster: A mariculture candidate for the Caribbean? *World Aquaculture* **22**:60-63.
- Phillips, B.F. and J.D. Booth. 1994. Design, use, and effectiveness of collectors for catching the puerulus stage of spiny lobsters. *Reviews in Fisheries Science* **2**:181-285.
- Reid, D. M., Corey, S., and M.H. Goodwin. 1991. An effective artificial habitat for collecting juvenile *Panulirus argus* in St. Kitts and Nevis, West Indies. *Proceedings of the Gulf and Caribbean Fisheries Institute* **44**:558-564.
- Ryther, J.H., W.A. Lellis, S.D. Bannerot, and J.A. Chaiton. [1988]. Spiny Lobster Mariculture. Final Report US Agency for International Development, Washington, D.C. USA. 42 pp.
- Witham, R., R.M. Ingle, and H.W. Sims. 1964. Notes on postlarvae of *Panulirus argus*. *Quarterly Journal of the Florida Academy of Sciences* **27**:289-297.